

Introduction

Example

Compute:

(a) $(1 + 3\sqrt{2}) + (-3 + 2\sqrt{2})$

(b) $(2 + 2\sqrt{2}) - (4 - \sqrt{2})$

(c) $(3 + \sqrt{2})(4 - 2\sqrt{2})$

(d) $\frac{1 + \sqrt{2}}{-3 + 2\sqrt{2}}$

Example

Suppose $x^2 = 2$, compute

(a) $(1 + 3x) + (-3 + 2x)$

(b) $(2 + 2x) - (4 - x)$

(c) $(3 + x)(4 - 2x)$

(d) $\frac{1 + x}{-3 + 2x}$

We introduce a new number, i , such that $i^2 = -1$, and write numbers as $a + ib$ where $a, b \in \mathbb{R}$.

Fact — All the normal rules of addition and multiplication still work

$$(1 + i) + (4 + 5i) = (1 + 4) + (1 + 5)i = 5 + 6i$$

$$\begin{aligned}(3 + 2i) \times (4 + 5i) &= 3 \times (4 + 5i) + 2i \times (4 + 5i) \\ &= 12 + 15i + 8i - 10 \\ &= 2 + 23i\end{aligned}$$

Example

Solve the equation $(2 + i)z = 3 + 4i$

Definition. If $z = a + ib \in \mathbb{C}$, $a, b \in \mathbb{R}$ then we call a the **real part** of z , $a = \operatorname{Re}(z)$ and b the **imaginary part** of z , $b = \operatorname{Im}(z)$.

Definition. If $z = a + ib \in \mathbb{C}$, $a, b \in \mathbb{R}$ is a complex number, then its complex conjugate $z^* = a - ib$

Example

Prove that:

(a) $(z + w)^* = z^* + w^*$

(b) $(zw)^* = z^*w^*$

(c) $(z^*)^* = z$

(d) $a^* = a$ if $a \in \mathbb{R}$

Example

Find the roots of $z^2 + 4z + 5 = 0$

Example


If f is a polynomial with *real* coefficients then its roots come in complex-conjugate pairs

Example

Solve $z^2 + i = 0$

Fact (Algebraic Closure) — If $f \in \mathbb{C}[X]$ is a polynomial with coefficients in \mathbb{C} then it has a complex root.

The Complex Plane



Definition. The **modulus** of a complex number, $z = a + bi \in \mathbb{C}, a, b \in \mathbb{R}$,

$$|z| = \sqrt{a^2 + b^2}$$

Definition. The **argument** of a complex number, $z = a + bi \in \mathbb{C}, a, b \in \mathbb{R}$, is the angle in *radians* measured anticlockwise from the positive real axis.

$$\arg(z) = \arctan\left(\frac{b}{a}\right) \quad (*)$$

$\arg(z) \in (-\pi, \pi]$ or $\arg(z) \in [0, 2\pi)$.

Example

Find the modulus and argument of $1 + i, 2i, -1 + 2i, -1 - i$.

Fact — If $z_1, z_2 \in \mathbb{C}$ then $|z_1 z_2| = |z_1| |z_2|$ and $\arg(z_1 z_2) = \arg(z_1) + \arg(z_2) \pmod{2\pi}$

Example

Find the complex number with modulus 2 and argument $\frac{\pi}{3}$

Definition. The **modulus-argument form** of a complex number is

$$z = r(\cos \theta + i \sin \theta) = r \operatorname{cis} \theta$$

Example

Show that $\sin(A + B) = \sin A \cos B + \cos A \sin B$ and find a similar formula for $\cos(A + B)$

Complex Geometry

Example

Show that $|z + w| \leq |z| + |w|$

Example


Sketch

(a) $|z - (1 + i)| = 2$

(b) $\arg(z - 1) = \frac{\pi}{3}$

(c) $\operatorname{Re}(z) = 3$


(d) $|z - 1| = |z - i|$



Example

Sketch

(a) $|z - 1| = 2|z - i|$



Actual Complex Geometry

Fact (Pro Tips) —

- Addition is translation
- Multiplication by a positive real number is enlargement
- Conjugation is reflection in the real-axis.
- Multiplication by $\text{cis}\theta$ is (anti-clockwise) rotation about the origin by θ

Example (Bottema's Theorem)

in any given triangle ABC , construct squares on any two adjacent sides, for example AC and BC . The midpoint of the line segment that connects the vertices of the squares opposite the common vertex, C , of the two sides of the triangle is independent of the location of C

Example (Napoleon's Theorem)

On the exterior of triangle ABC three new equilateral triangles $AC'B$, $BA'C$ and $CB'A$ are constructed. Prove that the centroids of these triangles are the vertices of an equilateral triangle