

## Approximating Curves by Polynomials

### Why do we care about polynomial approximations?

Polynomials are easy to work with - we can differentiate them, integrate them, and evaluate them using only basic arithmetic. But many important functions like  $e^x$ ,  $\sin x$ , and  $\ln(1+x)$  are not polynomials. If we could approximate these functions with polynomials, calculations would become much simpler.

#### Example

Find an approximation to  $f(x) = \sqrt{1+x}$  near  $x = 0$ .

**Definition** (Agreement to the  $n$ th degree). Two functions  $f$  and  $g$  agree to the  $n$ th degree at  $x = 0$  if:

$$f(0) = g(0), \quad f'(0) = g'(0), \quad f''(0) = g''(0), \quad \dots, \quad f^{(n)}(0) = g^{(n)}(0)$$

## The Maclaurin Polynomial Formula

### Theorem (Maclaurin Polynomial)

The polynomial of degree  $n$  which agrees with  $f(x)$  to the  $n$ th degree at  $x = 0$  is:

$$p_n(x) = f(0) + \frac{f'(0)}{1!}x + \frac{f''(0)}{2!}x^2 + \dots + \frac{f^{(n)}(0)}{n!}x^n = \sum_{r=0}^n \frac{f^{(r)}(0)}{r!}x^r$$

**Example**

Find the Maclaurin polynomial of degree 4 for  $f(x) = (1 + x)^{-3}$ .

**Example**

Find the Maclaurin polynomial of degree 4 for  $f(x) = \sin(2x)$ .

## From Polynomials to Series

**Definition** (Maclaurin Series). The **Maclaurin series** of  $f(x)$  is the infinite series:

$$f(x) = \sum_{r=0}^{\infty} \frac{f^{(r)}(0)}{r!} x^r = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \dots$$

[The set of  $x$  values for which this series converges to  $f(x)$  is called the **interval of validity**.]

## Standard Maclaurin Series

### The Exponential Function

**Example**

Find the Maclaurin series for  $f(x) = e^x$ .

### Trigonometric Functions

**Example**

Find the Maclaurin series for  $\sin x$

## The Natural Logarithm

**Example**

Find the Maclaurin series for  $f(x) = \ln(1 + x)$ .

Fact (Standard Maclaurin Series) —

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots + \frac{x^r}{r!} + \cdots \quad \text{for } x \in \mathbb{R}$$

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \cdots + (-1)^r \frac{x^{2r+1}}{(2r+1)!} + \cdots \quad \text{for } x \in \mathbb{R}$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \cdots + (-1)^r \frac{x^{2r}}{(2r)!} + \cdots \quad \text{for } x \in \mathbb{R}$$

$$\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \cdots + (-1)^{r-1} \frac{x^r}{r} + \cdots \quad \text{for } -1 < x \leq 1$$

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \cdots \quad \text{see below}$$

For  $(1+x)^n$ : valid for all  $x$  if  $n \in \mathbb{N}$ , otherwise for  $|x| < 1$ .

## Applications of Maclaurin Series

### Numerical Calculations

**Example**

Find  $e$  to 4 decimal places.

**Example (WJEC)** (a) Show that the first two non-zero terms in the Maclaurin expansion of  $\sin^{-1} x$  are given by

$$\sin^{-1} x = x + \frac{x^3}{6} + \dots$$

- (b) By writing  $x = \frac{1}{2}$ , deduce an approximation to  $\pi$  as a rational fraction in its lowest terms.
- (c) The equation  $\sin^{-1} x = 1.002x$  is satisfied by a small positive value of  $x$ . Find an approximation to this value, giving your answer correct to three decimal places.

## Composite Functions

**Example**

Find the Maclaurin expansion of  $e^{2x} \cos 3x$  up to the term in  $x^4$ .

**Tip**

When multiplying series, be systematic! Write out both series, then collect terms of the same degree. Remember that you only need terms up to the required power.

## Intervals of Validity

**Fact** — The interval of validity is the set of  $x$  values for which the Maclaurin series converges to the function value. Key points:

- For  $e^x$ ,  $\sin x$ ,  $\cos x$ : valid for all real  $x$
- For  $\ln(1 + x)$ : valid for  $-1 < x \leq 1$
- For  $(1 + x)^n$  with  $n \notin \mathbb{N}$ : valid for  $|x| < 1$
- When composing functions, check that the inner function stays within the validity range

### Example

For what values of  $x$  is the Maclaurin series for  $\ln(1 + 3x^2)$  valid?

**Example** (OCR June 2010 Q3)

Given that the first three terms of the Maclaurin series for  $(1 + \sin x)e^{2x}$  are identical to the first three terms of the binomial series for  $(1 + ax)^n$ , find the values of the constants  $a$  and  $n$