

## Revision

**Example (Algebra)**

Suppose  $z = 4 + i$ ,  $w = 3 - 4i$ . Find

- (a)  $z + w$
- (b)  $z - w$
- (c)  $zw$
- (d)  $\frac{z}{w}$
- (e)  $z^*$
- (f)  $|z|$  and  $|w|$
- (g)  $\arg(z)$  and  $\arg(w)$

**Example**

$$\text{Solve } z^2 + (-4 - 3i)z + (1 + 7i) = 0$$

**Complex Geometry****Example**

Show that  $|z + w| \leq |z| + |w|$

**Example**


Sketch

(a)  $|z - (1 + i)| = 2$

(b)  $\arg(z - 1) = \frac{\pi}{3}$

(c)  $\operatorname{Re}(z) = 3$


(d)  $|z - 1| = |z - i|$



**Example**

Sketch

(a)  $|z - 1| = 2|z - i|$



## Euler's Formula

**Example**

Calculate  $e^{ix}$  using the Maclaurin series

**Example**

What are:

(a)  $e^0$

(b)  $e^{i\pi}$

(c)  $e^{i\frac{\pi}{6}}$

(d)  $i^i$

**Definition.** A complex number is in **exponential form** when it is written as

$$z = re^{i\theta}, r \geq 0$$

**Fact** — We can now write complex numbers in 4 ways:

$$z = x + iy = re^{i\theta} = r \operatorname{cis} \theta = (r; \theta)$$

**Example**

Write  $1 + \sqrt{3}i$  in exponential form.

**Example**

Write  $5e^{i\frac{5\pi}{6}}$  in cartesian form.

**Example**

Suppose  $z_1 = r_1 e^{i\theta_1}$ ,  $z_2 = r_2 e^{i\theta_2}$  then calculate

(a)  $z_1 z_2$

(b)  $\frac{z_1}{z_2}$

(c)  $z_1^n$

**Example** (OCR Jan 2009 - FP3 Q2)

(i) Express  $\frac{\sqrt{3}+i}{\sqrt{3}-i}$  in the form  $re^{i\theta}$ , where  $r > 0$  and  $0 \leq \theta < 2\pi$ . [3]

(ii) Hence find the smallest positive value of  $n$  for which  $\left(\frac{\sqrt{3}+i}{\sqrt{3}-i}\right)^n$  is real and positive. [2]

## de Moivre's Theorem

Fact —


$$(\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta$$

**Example**

Write  $(\cos \frac{2\pi}{7} + i \sin \frac{2\pi}{7})^9$  in the form  $a + bi$


**Example**Compute  $\sqrt[3]{1+i}$ **Example**

Plot these values on an Argand diagram.



**Example**Solve  $z^n - 1 = 0$ **Example**

Plot these values on an Argand diagram.



## Trigonometry

**Example**

Express  $\sin 3\theta$  in terms of  $\sin \theta$ .

**Example** (OCR Jan 2010 FP3 Q7)

(i) Solve the equation  $\cos 6\theta = 0$ , for  $0 < \theta < \pi$ . [3]

(ii) By using de Moivre's theorem, show that

$$\cos 6\theta \equiv (2\cos^2\theta - 1)(16\cos^4\theta - 16\cos^2\theta + 1). \quad [5]$$

(iii) Hence find the exact value of

$$\cos\left(\frac{1}{12}\pi\right)\cos\left(\frac{5}{12}\pi\right)\cos\left(\frac{7}{12}\pi\right)\cos\left(\frac{11}{12}\pi\right),$$

justifying your answer. [5]

Fact —

$$\cos \theta = \operatorname{Re}\left(e^{i\theta}\right) = \frac{e^{i\theta} + e^{-i\theta}}{2}$$

$$\sin \theta = \operatorname{Im}\left(e^{i\theta}\right) = \frac{e^{i\theta} - e^{-i\theta}}{2i}$$

**Example**

Express  $\cos^6 \theta$  in terms of cos and sin of multiples of  $\theta$

**Example** (OCR June 2008 - FP3 Q4)

(i) By expressing  $\cos \theta$  in terms of  $e^{i\theta}$  and  $e^{-i\theta}$ , show that

$$\cos^5 \theta \equiv \frac{1}{16}(\cos 5\theta + 5 \cos 3\theta + 10 \cos \theta)$$

[5]

(ii) Hence solve the equation  $\cos 5\theta + 5 \cos 3\theta + 9 \cos \theta = 0$  for  $0 \leq \theta \leq \pi$ .

[4]

**Example**

Calculate

$$\sin \theta + \frac{1}{3} \sin 3\theta + \frac{1}{9} \sin 5\theta + \frac{1}{27} \sin 7\theta + \dots$$

## Osborn's Rule

**Fact (Osborn's Rule)** — Any trigonometric identity can be converted to a hyperbolic identity by:

- (1) Replacing  $\cos \theta$  with  $\cosh \theta$
- (2) Replacing  $\sin \theta$  with  $\sinh \theta$
- (3) Negating any term containing a product of two sines (e.g.  $\sin^2 \theta$ ,  $\sin A \sin B$ ,  $\tan^2 \theta$ )

**Remark** (Why does Osborn's rule work?). Substituting  $i\theta$  into Euler's formula gives us a connection between trig and hyperbolic functions:

$$\begin{aligned}\cos(i\theta) &= \frac{e^{i(i\theta)} + e^{-i(i\theta)}}{2} = \frac{e^{-\theta} + e^{\theta}}{2} = \cosh \theta \\ \sin(i\theta) &= \frac{e^{i(i\theta)} - e^{-i(i\theta)}}{2i} = \frac{e^{-\theta} - e^{\theta}}{2i} = \frac{-(e^{\theta} - e^{-\theta})}{2i} = \frac{i(e^{\theta} - e^{-\theta})}{2} = i \sinh \theta\end{aligned}$$

So if we take any trig identity and substitute  $\theta \rightarrow i\theta$ , every cos becomes cosh and every sin becomes  $i \sinh$ . When two sines are multiplied together, we get  $(i \sinh)(i \sinh) = -\sinh^2$ , which explains the sign change.

### Example

Convert the following identities to their hyperbolic equivalents using Osborn's rule:

- (a)  $\cos^2 \theta + \sin^2 \theta = 1$
- (b)  $\sin 2\theta = 2 \sin \theta \cos \theta$
- (c)  $\cos 2\theta = \cos^2 \theta - \sin^2 \theta$
- (d)  $1 + \tan^2 \theta = \sec^2 \theta$

### Example

Use Osborn's rule to find a formula for  $\cosh 3\theta$  in terms of  $\cosh \theta$ .

## Fibonometry

The Fibonacci numbers  $F_n$  and Lucas numbers  $L_n$  are defined by the same recurrence  $X_{n+1} = X_n + X_{n-1}$ , with initial conditions:

$$F_0 = 0, F_1 = 1 \qquad L_0 = 2, L_1 = 1$$

**Fact (Binet's Formulas)** — Let  $\phi = \frac{1+\sqrt{5}}{2}$  (the golden ratio) and  $\psi = \frac{1-\sqrt{5}}{2}$ . Then:

$$F_n = \frac{\phi^n - \psi^n}{\sqrt{5}} \qquad L_n = \phi^n + \psi^n$$

**Remark.** Compare Binet's formulas with the exponential forms of sine and cosine:

$$\sin \theta = \frac{e^{i\theta} - e^{-i\theta}}{2i} \qquad \cos \theta = \frac{e^{i\theta} + e^{-i\theta}}{2}$$

The Fibonacci numbers  $F_n$  play the role of sin, and the Lucas numbers  $L_n$  play the role of cos. This analogy, which Conway called **Fibonometry**, lets us translate trigonometric identities into Fibonacci/Lucas identities.

The following identities are parallel:

Trigonometric Identity	Fibonacci/Lucas Identity
$\sin(A + B) + \sin(A - B) = 2 \sin A \cos B$	$F_{m+n} + F_{m-n} = F_m L_n$ (for $n$ even)
$\sin(A + B) - \sin(A - B) = 2 \cos A \sin B$	$F_{m+n} - F_{m-n} = F_n L_m$ (for $n$ even)
$\cos(A + B) + \cos(A - B) = 2 \cos A \cos B$	$L_{m+n} + L_{m-n} = L_m L_n$ (for $n$ even)
$\cos(A + B) - \cos(A - B) = -2 \sin A \sin B$	$L_{m+n} - L_{m-n} = 5F_m F_n$ (for $n$ even)
$\cos^2 \theta - \sin^2 \theta = \cos 2\theta$	$L_n^2 - 5F_n^2 = 4(-1)^n$
$\sin 2\theta = 2 \sin \theta \cos \theta$	$F_{2n} = F_n L_n$

**Remark.** When  $n$  is odd, the + and - signs swap in the Fibonacci/Lucas identities (due to  $\psi^n$  being negative for odd  $n$ ).

### Example

Verify that  $F_{2n} = F_n L_n$  using Binet's formulas.

**Example**

Use Fibonometry to prove that  $L_n^2 - 5F_n^2 = 4(-1)^n$ .