

Translations


Throughout this topic, $y = f(x)$ is a known graph and we ask what happens to it when the equation is altered. The base curve here is $y = x^3$.

Fact —

$y = f(x) + a$ translation by $\begin{pmatrix} 0 \\ a \end{pmatrix}$ (y -values all increase by a)

$y = f(x + a)$ translation by $\begin{pmatrix} -a \\ 0 \end{pmatrix}$ (the input $x + a$ reaches each value *earlier*)

Changes *outside* f act on y , in the expected direction. Changes *inside* f act on x , in the opposite direction.



Example

The point $P(3, 7)$ lies on $y = f(x)$. Write down the image of P on:

1. $y = f(x) - 5$
2. $y = f(x + 2)$
3. $y = f(x - 1) + 4$

Example

The graph of $y = x^2$ is translated by $\begin{pmatrix} 3 \\ -2 \end{pmatrix}$. Find the equation of the image in the form $y = x^2 + bx + c$.

Textbook Exercises: SPS Course 3.2, Exercise 2 Q1–6

Stretches and Reflections

Fact —	$y = af(x)$	stretch, scale factor a , parallel to the y -axis
	$y = f(ax)$	stretch, scale factor $\frac{1}{a}$, parallel to the x -axis
	$y = -f(x)$	reflection in the x -axis
	$y = f(-x)$	reflection in the y -axis

The inside/outside rule again: inside acts on x , opposite to expectation.

Example


The point $Q(6, -2)$ lies on $y = f(x)$. Write down the image of Q on:

1. $y = 3f(x)$
2. $y = f(2x)$
3. $y = f(-x)$
4. $y = -f(x)$

Trigonometric Graphs

Example

On the same axes for $0^\circ \leq x \leq 360^\circ$, sketch $y = \sin x$, $y = 2 \sin x$ and $y = \sin 2x$. State the number of solutions of $\sin 2x = \frac{1}{2}$ in this interval.



Example

Describe the transformation taking $y = \cos x$ to:

1. $y = \cos x - 3$
2. $y = \cos\left(\frac{x}{2}\right)$
3. $y = -\cos x$

Textbook Exercises: SPS Course 3.2, Exercise 2 Q7–12

Recognising Transformations


Example

In each case, describe a single transformation taking the first graph to the second:

1. $y = x^2$ to $y = x^2 + 6x + 9$
2. $y = \frac{1}{x}$ to $y = \frac{3}{x}$
3. $y = 2^x$ to $y = 2^{x+1}$ — find a second, different, single transformation that also works.

Example

The diagram shows $y = f(x)$, which has a maximum at $A(2, 3)$ and crosses the axes at $(0, 1)$ and $(5, 0)$.



Sketch the following, marking the images of all three points:

1. $y = f(x - 3)$
2. $y = 2f(x)$
3. $y = -f(x)$
4. $y = f(-x)$

Example

The curve $y = x^3$ is reflected in the x -axis, then translated by $\begin{pmatrix} 0 \\ 4 \end{pmatrix}$. Find the equation of the result, and the coordinates of the point where it crosses the x -axis.

Textbook Exercises: SPS Course 3.2, Exercise 2 (remaining questions) and Revision Exercise 3.2